

Quantum spin liquids of Rydberg excitations in a honeycomb lattice induced by density-dependent Peierls phases

Simon Ohler¹, Maximilian Kiefer-Emmanouilidis,^{1,2} and Michael Fleischhauer¹

¹*Department of Physics and Research Center OPTIMAS, University of Kaiserslautern, D-67663 Kaiserslautern, Germany*

²*German Research Centre for Artificial Intelligence, Embedded Intelligence, D-67663 Kaiserslautern, Germany*



(Received 21 February 2022; accepted 5 January 2023; published 28 February 2023)

We show that the nonlinear transport of bosonic excitations in a two-dimensional honeycomb lattice of spin-orbit-coupled Rydberg atoms gives rise to disordered quantum phases which are topological and may be candidates for quantum spin liquids. As recently demonstrated by Lienhard *et al.* [*Phys. Rev. X* **10**, 021031 (2020)] the spin-orbit coupling breaks time-reversal and chiral symmetries and leads to a tunable density-dependent complex hopping of spin excitations which behave as hard-core bosons. Using exact diagonalization (ED), we numerically investigate the phase diagram resulting from the competition between density-dependent and direct transport terms as well as density-density interactions. In mean-field approximation there is a phase transition from a condensate to a 120° phase when the amplitude of the complex hopping exceeds that of the direct one. In the full model a new phase emerges close to the mean-field critical point as a result of quantum correlations induced by the density dependence of the complex hopping. We show that without density-density interactions this phase is a genuine disordered one, has large spin chirality, and is characterized by a nontrivial many-body Chern number. The Chern number is found to be robust to disorder. ED simulations of small lattices with up to 30 lattice sites give indications for a nondegenerate ground state with finite spin and collective gaps and thus for a bosonic integer quantum Hall phase, protected by $U(1)$ symmetry. On the other hand, while staying finite, the many-body gap varies substantially when different twisted boundary conditions are applied, which points to a gapless phase. For very strong negative nonlinear hopping amplitudes we find another disordered regime with vanishing spin gap. This phase also has a large spin chirality and could be a gapless spin liquid but lies outside the parameter regime experimentally accessible in the Rydberg system.

DOI: [10.1103/PhysRevResearch.5.013157](https://doi.org/10.1103/PhysRevResearch.5.013157)

I. INTRODUCTION

Over recent years, Rydberg atoms have become a versatile and robust platform to explore many-body quantum spin physics in the regime of strong correlations [1–6] and for quantum information processing [7–11]. Using high principal quantum numbers, Rydberg-excited atoms have sizable interactions even at distances of several micrometers, while their lifetime is on the order of milliseconds. These properties make Rydberg atoms especially well suited to explore many-body quantum phenomena such as the recently experimentally realized symmetry-protected topological phases [12] and quantum dimer models [13].

Particularly interesting and still poorly understood many-body phases of spin systems are those where zero-point quantum fluctuations prevent magnetic order of any kind, which often requires frustration. The possibility of such liquid ground states was first pointed out by Anderson in his seminal 1973 work on antiferromagnets [14,15]. Since then, there has

been an ongoing search for experimental realizations of such a quantum spin liquid (QSL) [16–23]. The QSL may be gapped or gapless and in the first case can be topological, where the topological order either can be protected by symmetries associated with short-range entanglement [24,25] or can be intrinsic, in which case the state is long-range entangled [26,27]. Despite the decades-long interest in quantum spin liquids, their clear identification in realistic materials remains a major challenge due to the scarcity of highly entangled states in real solid-state materials and the lack of simple experimental signatures of spin liquids [28–30]. Thus it is natural to ask whether QSLs can be realized in experimentally accessible model systems such as arrays of Rydberg atoms. First signatures of a QSL have indeed been found in such a model system in a recent breakthrough experiment with Rydberg atoms in an artificially assembled two-dimensional array of microtraps [13]. In that experiment, ground and Rydberg states of the atom form a spin- $\frac{1}{2}$ system. The atoms were placed on the links of a kagome lattice and driven under conditions of Rydberg blockade [8], which effectively realizes a quantum dimer model, for which a QSL ground state has been predicted [31,32]. The dimer states are formed by three nearby atoms, out of which at most one can be excited due to Rydberg blockade.

Inspired by recent experimental work [33], we here propose and analyze a lattice spin model, based on Rydberg

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.




FIG. 1. Honeycomb lattice with a two-site unit cell (sublattices A and B) of trapped atoms excited to two different Rydberg states $|1\rangle$ and $|0\rangle$, forming spin- $\frac{1}{2}$ systems. As indicated, spin-orbit coupling induced by an external magnetic field leads to nonlinear, complex (chiral) second-order hopping processes to the next-nearest-neighbor (NNN) hopping in addition to direct nearest-neighbor (NN) hopping. The relevant level structure of a single atom is shown in the inset. The NNN hopping is facilitated by virtual transitions from $|0\rangle$ to the off-resonant state $|+\rangle$ and can be controlled by varying its detuning Δ .

atoms on a honeycomb lattice as shown in Fig. 1, where the competition between nearest and next-nearest XY spin couplings leads to frustration. Different from Ref. [13] the spin degree of freedom is formed by two Rydberg states of the atoms. The spin can hop from one lattice site to the next by dipolar exchange interactions. As shown in Ref. [33], in such systems spin-orbit coupling induced by an external magnetic field explicitly breaks time-reversal and chiral symmetry and leads to a density-dependent, second-order complex hopping of excitations [34] which competes with the direct hopping. The strength of the complex hopping can be modified by tuning the energy separation between the Rydberg states. Different from Ref. [13] the Hamiltonian describing the system conserves the number of excitations, i.e., it has $U(1)$ symmetry, and we here consider half filling. We study the system using numerical simulations performing exact diagonalization (ED) on small lattices with up to 30 lattice sites with periodic boundary conditions using different cluster shapes.

If the effects of quantum fluctuations on the complex hopping are neglected, i.e., in mean-field approximation, there is a competition between the nearest-neighbor (NN) hopping, which tries to establish a condensate of the hard-core bosons, and the next-nearest-neighbor (NNN) hopping, driving the system into a 120° or spiral spin phase. A transition between the two phases occurs if the strength of the mean-field second-order hopping becomes comparable to the direct one.

In the full model we identify two new phases. In the vicinity of the mean-field critical point an intermediate phase emerges, which is bare of any simple spin order, has a nonvanishing spin chirality, and is characterized by a non-trivial many-body Chern number $C = 1$. Thus this phase is a candidate for a topological QSL. Its precise nature is, however, not completely clear and needs further investigation [35]. ED simulations for finite systems point to a gapped, nondegenerate ground state. The many-body gap does, however, vary in magnitude substantially when different twisted

boundary conditions are applied. A gapped and nondegenerate ground state would indicate symmetry-protected topological (SPT) order. On the other hand, the Chern number of $C = 1$, which is robust to potential disorder, is odd and thus different from the even values found in other systems showing a symmetry-protected bosonic integer quantum Hall (BIQH) effect [24,36–39] and expected from general classification arguments [24]. An odd value of the Chern number would, on the other hand, be consistent with a gapless QSL. In this case the collective gap should vanish in the thermodynamic limit and the ground state should become degenerate.

When the second-order hoppings have the opposite sign and become strong, another disordered spin phase emerges, which is again chiral but gapless. We believe that this phase is a candidate for a gapless chiral spin liquid but the parameter regime is outside of what can be realized in Rydberg systems.

Hard-core bosons on a honeycomb lattice with frustrated next-nearest-neighbor hopping were studied in Refs. [40,41], where the authors reported evidence for a particular type of gapless spin liquids, called a Bose metal, i.e., a QSL with a “Fermi-like” surface in momentum space. Subsequent density matrix renormalization group (DMRG) simulations, however, showed weak density order [42], which demonstrates that an unambiguous identification of the nature of spin liquids using ED simulations is difficult.

The outline of this paper is as follows: In Sec. II we introduce the many-body Hamiltonian of Rydberg spin excitations in a two-dimensional honeycomb array of trapped atoms. An overview of the ground-state phases is given in Sec. III, where we show that the presence of complex and density-dependent second-order hopping processes gives rise to two disordered spin phases in addition to a trivial Bose-Einstein condensate (BEC) phase and a spiral, or 120° , phase present in a mean-field Hamiltonian. We comment on the possible nature of the QSL in Sec. IV and discuss the effects of longer-range interactions in Sec. V. Finally, a summary and discussion of the results are given in Sec. VI.

II. MODEL FOR RYDBERG EXCITATIONS ON A HONEYCOMB LATTICE

We consider a honeycomb array of microtraps filled with one atom each as shown in Fig. 1. Each site has three nearest neighbors (NNs) of the opposite sublattice and six next-nearest neighbors (NNNs) of the same sublattice A and B, respectively. As has been demonstrated in recent works [3,43–45], a deterministic and defect-free preparation of such lattice structures with characteristic separations in the micrometer scale is possible and state of the art. Each atom is excited into high-lying Rydberg states, e.g., within the $60S_{1/2}$ and $60P_{3/2}$ manifold of ^{87}Rb as in Ref. [33]. Application of an external magnetic field perpendicular to the plane leads to a level structure where only three magnetic sublevels are relevant, indicated in Fig. 1. Two of them, here labeled as $|0\rangle$ and $|1\rangle$, form an effective spin- $\frac{1}{2}$ system, and we are interested in the many-body dynamics of these spins. Dipolar coupling between the Rydberg-excited atoms leads to a hopping of spin excitations (XY coupling) with an amplitude J proportional to $1/r^3$, with r being the atom separation. In the case of a transition between the $m_J = \pm\frac{1}{2}$ sublevels of $S_{1/2}$ and $P_{3/2}$, as

mentioned above, $J = d^2/(8\pi\epsilon_0 r^3)$, with d being the dipole matrix element between states $|0\rangle$ and $|1\rangle$. The third level, denoted by $|+\rangle$, is used to facilitate a second-order, off-resonant spin exchange process that is associated with a geometry-dependent complex phase (see Fig. 1) and which depends on the spin state of the intermediate atom. The microscopic physics of the system has been studied both theoretically and experimentally in a minimal setup [33,46], and the relevant terms of the many-body Hamiltonian have been introduced and studied for a different lattice in Ref. [47]. For a detailed derivation of the Hamiltonian we thus refer to these publications. Following along the same lines as in Ref. [47], we write down the effective Hamiltonian for Rydberg excitations in the hard-core boson language (we use $\hbar = 1$):

$$\hat{H} = -J \sum_{\langle i,j \rangle} \hat{b}_j^\dagger \hat{b}_i - 2gJ \sum_{\langle\langle i,j \rangle\rangle} \hat{b}_j^\dagger \hat{b}_i e^{\pm \frac{2\pi i}{3}} (1 - \hat{n}_{ij}) + 2gJ \sum_{(i,j)} \hat{n}_i \hat{n}_j, \quad (1)$$

where $e^{\pm \frac{2\pi i}{3}} = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$ and $\hat{b}_i^\dagger, \hat{b}_i$ create or destroy a hard-core boson on site i , respectively. $\langle i, j \rangle$ and $\langle\langle i, j \rangle\rangle$ refer to NNs and NNNs, where both $i \rightarrow j$ and $j \rightarrow i$ are included in the sum. The sign of the complex phase as well as the intermediate site of the NNN hopping terms connecting sites i and j is indicated in Fig. 1 by differently colored and bent arrows, respectively. Thus the hopping between two nearest neighbors i and j of the same sublattice is controlled by a site of the opposite sublattice, located in between the two and with particle number \hat{n}_{ij} . Note that the complex phase picked up in a closed loop around a honeycomb plaquette corresponds to exactly one flux quantum. Furthermore, in the nearest-neighbor interaction term we have assumed conservation of particle number and dropped the constant energy shift

$$\sum_{(i,j)} \hat{n}_i (1 - \hat{n}_j) \rightarrow - \sum_{(i,j)} \hat{n}_i \hat{n}_j. \quad (2)$$

All processes contained in the Hamiltonian are shown in Fig. 1. This includes (1) NN hopping with constant amplitude J which depends on the atomic level structure and the spatial separation between the atoms, (2) NNN hopping that is density dependent, possesses a staggered complex phase, and scales with an additional parameter g , and (3) NN interaction that also scales with g . Terms connecting sites further apart are smaller in magnitude and will be neglected in first approximation. We will discuss their influence at the end of the paper. In Eq. (1) the strength of the nonresonant processes g is given by $g = 27J/(2\Delta)$, where Δ denotes the detuning between two Rydberg states of the atoms. The factor 27 stems from Clebsch-Gordan coefficients and factors in the microscopic Hamiltonian (see Ref. [47]). The additional factor $\frac{1}{2}$ in the definition of g is introduced to be consistent with Ref. [47], where the same atomic setup is studied on a zigzag chain. Most importantly, the magnitude and sign of g can be controlled by the detuning of the internal state $|+\rangle$. In order to be able to neglect the population of the off-resonant state, the detuning cannot be too small, i.e., $J/|\Delta| \ll 1$, but values of $|g| \sim 2$ are possible. In this paper we consider half filling of hard-core bosons corresponding to a vanishing total magnetization.




FIG. 2. Collections of cluster shapes and sizes used for numerical calculations. The nomenclature is consistent with Ref. [41].

Let us first discuss some general aspects of Hamiltonian (1). The presence of complex hopping amplitudes means that time-reversal symmetry is explicitly broken. The microscopic origin of this is the magnetic field used to select the specific sublevels of the Rydberg atoms. Secondly, without the nonlinear term in the NNN hopping the model is symmetric under a combined time-reversal and particle-hole transformation at half filling, but this symmetry is broken by the term $(1 - \hat{n}_{ij})$. The NN density-density interaction corresponding to a ferromagnetic ($gJ < 0$) or antiferromagnetic ($gJ > 0$) Ising term would drive the system into a density-ordered state. The NNN hopping which is of the same strength, however, prevents the formation of a state with ferromagnetic or antiferromagnetic density order. Therefore the possible phases are essentially governed by the competition of the NN and NNN hopping terms and the action of the nonlinear term in the NNN hopping amplitude.

III. GROUND-STATE PHASES AND EFFECTS OF NONLINEAR HOPPING

In order to investigate the different ground-state phases of the model (1), we use exact diagonalization (ED) on finite lattices using periodic (or twisted) boundary conditions. In order to reduce boundary effects, we perform calculations on hexagonal clusters of varying shapes and sizes. The clusters that we use are shown in Fig. 2. Using the Lanczos algorithm [48], we gain access to the ground-state wave function.

To obtain a general overview of the phase diagram as a function of the parameter g , we consider the change in the ground-state wave function $|\Phi_0\rangle$ upon infinitesimal changes in g . We can quantify this by the dimensionless, intensive fidelity metric

$$f(g) = \frac{2}{L} \frac{1 - |\langle \Phi_0(g) | \Phi_0(g + \delta g) \rangle|}{(\delta g)^2}, \quad \delta g \rightarrow 0. \quad (3)$$

Here, L represents the number of sites in the system. This quantity has been shown to be a useful indicator of quantum phase transitions [49] and has since been used in numerous condensed-matter applications [40,41,50,51]. By computing the overlap of the ground-state wave function with itself under small changes in g , we are able to detect the regions in parameter space where the system's ground state changes rapidly, indicating a possible quantum phase transition (QPT). In a finite system this quantity will always be finite and not show a Dirac- δ -like behavior, which we expect at the critical point of a QPT in an infinite system. Therefore we can use it only as a rough guide to separate different parameter regimes.




FIG. 3. Ground-state fidelity metric f for the mean-field system given in (5) as a function of the parameter g . The peak of f agrees well for the different system shapes and indicates a phase transition. For $g \lesssim 0.4$ we find a superfluid state, while for $g \gtrsim 0.4$ the system shows 120° order. The inset shows the region around the phase transition in more detail. Note that the curves for shapes 24C and 24D are virtually indistinguishable.

As we will show, the density-dependent hopping in (1) has a profound impact on the behavior of the system.

A. Mean-field dynamics: Competition between NN and NNN hopping

To gain some insight into the main competing terms in the Hamiltonian, we first study a mean-field approximation. To this end, we drop the nearest-neighbor interaction term and replace the density dependence of the NNN hopping term with a constant expectation value

$$(1 - \hat{n}_{ij}) \rightarrow (1 - \bar{n}). \quad (4)$$

Here, \bar{n} denotes the average density of the lattice. Since we consider half filling, we set $\bar{n} = 0.5$. The modified approximate Hamiltonian then reads

$$\hat{H}_{\text{MF}} = -J \sum_{\langle i,j \rangle} \hat{b}_j^\dagger \hat{b}_i - gJ \sum_{\langle\langle i,j \rangle\rangle} \hat{b}_j^\dagger \hat{b}_i e^{\pm \frac{2\pi}{3}i}. \quad (5)$$

This mean-field Hamiltonian is that of the Bose-Hubbard Haldane model. The corresponding single-particle band structure is that of Haldane's generalization of graphene in the topologically nontrivial regime [52]. However, as shown in Ref. [53], the hard-core boson character of the Rydberg excitations makes the many-body ground state topologically trivial. Note that while in one dimension hard-core bosons and fermions can behave identically under certain circumstances, this is decidedly not the case in two dimensions, and the generalization of the Wigner-Jordan transformation to two dimensions requires the introduction of effective gauge fields [54]. In Fig. 3 we have plotted the fidelity metric of the ground state as a function of the interaction strength g , using the mean-field Hamiltonian (5) with periodic boundary conditions (PBCs) on a torus. From the fidelity we see two regimes separated by a single peak. The left region is continuously connected to the trivial limit $g = 0$, where the system is in a BEC state. In order to understand the phase in the right region, it is sufficient to consider the case $g \gg 1$, where the NN hopping term is

irrelevant:

$$\hat{H}_{g \rightarrow \infty} = -gJ \sum_{\langle\langle i,j \rangle\rangle} \hat{b}_j^\dagger \hat{b}_i e^{\pm \frac{2\pi}{3}i}. \quad (6)$$

Now, the two triangular sublattices A and B of the hexagonal lattice are disconnected, and the internal dynamics in each of the sublattices is determined by the NNN hopping. Furthermore, we can rewrite the Hamiltonian in terms of spin- $\frac{1}{2}$ matrices

$$\hat{b}^\dagger \rightarrow \hat{S}^+ = \hat{S}^x + i\hat{S}^y, \quad \hat{b} \rightarrow \hat{S}^- = \hat{S}^x - i\hat{S}^y, \quad (7)$$

such that the NNN hopping term reads

$$\hat{H}_{g \rightarrow \infty} = -\frac{gJ}{2} \sum_{\Delta} (\mathbf{S}_{\Delta,1}^T D \mathbf{S}_{\Delta,2} + \mathbf{S}_{\Delta,2}^T D \mathbf{S}_{\Delta,3} + \mathbf{S}_{\Delta,3}^T D \mathbf{S}_{\Delta,1}). \quad (8)$$

Here, the vector operators $\mathbf{S}_{\Delta,i}$ are projections of the spins to the xy plane, the index Δ runs over all triangles of both triangular sublattices, and the index 1,2,3 iterates through a single triangle as indicated in the inset of Fig. 3. The matrix $D = D(2\pi/3)$ is the rotational matrix around the z axis with rotation angle of $2\pi/3$. With $D^3 = 1$ we can write $\hat{H}_{g \rightarrow \infty}$ in its final form:

$$\hat{H}_{g \rightarrow \infty} = -\frac{gJ}{2} \sum_{\Delta} (\mathbf{S}_{\Delta} + D \mathbf{S}_{\Delta,2} + D^2 \mathbf{S}_{\Delta,3})^2 + \text{const.} \quad (9)$$

For $g > 0$ we see that in the ground state the rotated spins $\mathbf{S}_{\Delta,1}$, $D \mathbf{S}_{\Delta,2}$, and $D^2 \mathbf{S}_{\Delta,3}$ need to be parallel, i.e., the spin vectors themselves have to be at an angle of 120° in the xy plane. We can confirm this calculation by considering the in-plane spin correlations, which we define as

$$C(\theta) = 4 \langle \hat{S}_i^{(0)} \hat{S}_j^{(\theta)} \rangle, \quad (10)$$

where

$$\hat{S}_j^{(\theta)} = \cos(\theta) \hat{S}_j^x + \sin(\theta) \hat{S}_j^y. \quad (11)$$

This correlation function detects whether both spin vectors are separated by an angle θ in the equatorial plane. The factor of 4 is introduced to normalize $C(\theta)$ to unity in the case of a perfectly correlated state.

In Fig. 4 we show the results for the in-plane spin correlations in the mean-field limit, which show a maximum at $\theta = 120^\circ$ for large g . The case of $g < 0$ can be understood similarly: Here, the sum over the rotated spin vectors has to vanish, which requires the spins to align in a parallel way in the xy plane. The $\theta = 120^\circ$ phase has a remaining $\text{SO}(2)$ symmetry. The lack of any correlation between NNs in the 120° phase for large g is a trivial result of the disconnected sublattices in this limit. The small deviation of $C(120^\circ)$ from the maximum value of $\frac{1}{2}$ [taking into account the $\text{SO}(2)$ symmetry] can be attributed to finite-size effects.

B. Full Hamiltonian

After having discussed the mean-field Hamiltonian, we now turn to the microscopically motivated full Hamiltonian (1), which includes a density-dependent, complex NNN hopping term and NN density-density interaction. In this case, the ground-state fidelity (3) is modified substantially, as shown




FIG. 4. In-plane spin-spin correlation (10) in mean-field approximation (5). (a) shows $C(\theta)$ for i and j being NNs (different sublattices); (b) depicts the NNN (same sublattice) correlation. All calculations were performed with periodic boundary conditions on shape 24C. The 120° regime as well as the parallel alignment in the BEC phase can be clearly seen. Furthermore, one recognizes that NNs are completely uncorrelated in the 120° phase.

in Fig. 5. Where for the mean-field model we had seen only one phase transition at g close to 0.4, we now see two sharp peaks of the fidelity in the vicinity of the mean-field critical point. Thus a new phase emerges in between the BEC ($g \rightarrow 0$) and the 120° order ($g \gg 0.4$), which we refer to as regime (or phase) II. Additionally, for $g \approx -5$ we see a behavior that is qualitatively different from the mean-field case. Due to the lack of a clear peak in that region it is unclear whether this indicates another true phase transition or a crossover. This area of the parameter space we denote as regime (or phase) I. We note, however, that in this region the condition $J/|\Delta| \ll 1$ resulting from microscopic physics of Rydberg interactions and required to neglect the population in level $|+\rangle$ is no longer fulfilled.

In the following we will characterize the new phases by different observables.

1. In-plane spin orientation

In Fig. 6 we show the results of the in-plane spin correlations for the full Hamiltonian (1). In the BEC as well as in the




FIG. 5. Ground-state fidelity metric f as a function of the parameter g for the full Hamiltonian (1). The peaks of f agree well for the different shapes and indicate potential phase transitions. The inset shows regime II in detail. The BEC and 120° -order regimes agree with the mean-field model, but two new regimes appear, which we label as regimes (or phases) I and II.




FIG. 6. (a) and (b) In-plane spin-spin correlation (10) as in Fig. 4, but for the full Hamiltonian (1). One recognizes a very similar behavior for the BEC and 120° phases as in the mean-field case. In phases I and II the in-plane spin correlations are suppressed.

120° phase the correlations are almost identical to the mean-field case, supporting our interpretation of these phases. In the narrow intermediate phase II the in-plane spin correlations are suppressed, and they are also reduced when entering and getting deeper into phase I. As opposed to the transition points to phase II the changes in the correlations at the transition into phase I are not sharp.

2. Spin chirality

The Peierls phases in the NNN hopping terms explicitly break time-reversal symmetry. Due to the absence of a mass term shifting the energy of sublattice A relative to that of sublattice B, the mean-field Hamiltonian, Eq. (5), preserves chiral symmetry, which amounts to a combination of time-reversal and particle-hole transformation. The nonlinear term in the complex NNN hopping amplitude of the full model, Eq. (1), however, breaks the chiral symmetry. Thus we expect that the disordered phases I and II are characterized by a significant spin chirality. The latter is defined as [55]

$$\chi = \langle \hat{\sigma}_i \cdot (\hat{\sigma}_j \times \hat{\sigma}_k) \rangle, \quad (12)$$

where $\hat{\sigma}$ is the three-component vector of Pauli operators in the spin- $\frac{1}{2}$ representation of the hard-core boson model. The indices $\{i, j, k\}$ in (12) are labeled in counterclockwise order around the elementary triangles of the honeycomb lattice as displayed in Fig. 7. Chiral symmetry would enforce $\chi = 0$. In Fig. 7 we plot the spin chirality on the three types of triangles as a function of the interaction strength g . We observe that in both disordered regimes their values are much larger than in the BEC and 120° -order phases. In phase I in Fig. 5, all three χ_i behave similarly, whereas in regime II their values differ in sign, indicating that the mediated interactions between the sublattices play an important role in the physics of regime II.

3. Spin order

a. Spin structure factor. In order to investigate the presence or absence of spin order in phases I and II, we now consider the spin structure factor, defined as

$$S(\mathbf{k}) = \frac{1}{L} \sum_{i,j=1}^N e^{-ik \cdot (r_i - r_j)} \langle \hat{S}_i \cdot \hat{S}_j \rangle, \quad (13)$$




FIG. 7. Visualization of the spin chirality measured for the full Hamiltonian (1). Top: We denote three different types of triangles by χ_i , $i \in \{1, 2, 3\}$. Bottom: Calculation of the spin chirality for the triangles displayed above. We observe that in regime I, all three chiralities behave similarly, while in regime II the values differ in sign.

where $\hat{\mathbf{S}}_j = (\hat{S}_j^x, \hat{S}_j^y, \hat{S}_j^z)^T$ is the full three-dimensional (3D) spin vector and L is the system size. In order to minimize the impact of finite-size effects, it is useful to impose a more general type of boundary conditions. Specifically, we consider twisted boundary conditions, where a particle picks up a phase upon hopping over the boundary. Therefore we average $S(\mathbf{k})$ over all low-energy configurations with twisted boundary conditions (see Sec. III B 3 c).

In Fig. 8 we show the results for the spin order, Eq. (13), for $L = 24$ (shape 24C). In the BEC regime ($g = 0$) the plot shows a single prominent peak around zero momentum, consistent with parallel spins in the xy plane as discussed before.




FIG. 8. Averaged spin structure factor over all low-energy configurations for (a) $g = -5.0$, (b) $g = 0$, (c) $g = 0.73$, and (d) $g = 4.0$. For $g = 4.0$, we see the well-known result for a 120° order, which is explained in the text. For $g = -5.0$ and $g = 0.73$ we see no prominent features after the averaging.




FIG. 9. Dimerization $D_{ij,kl}$ as defined in (14) for $g = 0.73$ (phase II). The linewidth and color encode the magnitude and sign, respectively. In this figure, we keep the reference bond i, j constant and vary k, l . The reference bond color box is shown with rounded corners.

The 120° order occurs for $g > 1$ such that NNN hopping terms and the density-density repulsion are dominant, whereas NN hopping is weak. Consequently, we can imagine the honeycomb lattice as two separate triangular sublattices. In this limit, the nearest-neighbor spin orientation in the xy plane is completely uncorrelated, while NNNs align themselves at an angle of $\frac{2\pi}{3} = 120^\circ$. This is demonstrated in the hexagonal peaks of Fig. 8(d), which are a well-known indicator of 120° order [56] (also referred to as spiral order). Additionally, as we have seen already in Fig. 6, we find that the in-plane spin correlation (10) vanishes for spins of different sublattices and prefers the angle $2\pi/3$ for the same sublattice.

b. Dimerization. In order to test for the presence of more involved orderings, we investigate whether a regular pattern of dimerized spins occurs in the system. To this end, we calculate the dimer-dimer correlation function introduced in Ref. [57]

$$D_{ij,kl} = \left\langle \left(\hat{\mathbf{S}}_i \cdot \hat{\mathbf{S}}_j - \frac{1}{4} \right) \left(\hat{\mathbf{S}}_k \cdot \hat{\mathbf{S}}_l - \frac{1}{4} \right) \right\rangle - \left\langle \hat{\mathbf{S}}_i \cdot \hat{\mathbf{S}}_j - \frac{1}{4} \right\rangle \left\langle \hat{\mathbf{S}}_k \cdot \hat{\mathbf{S}}_l - \frac{1}{4} \right\rangle, \quad (14)$$

where i, j and k, l are each NNs. The results of the calculation are shown in Fig. 9, where we observe only short-ranged correlations.

c. Randomly twisted boundary conditions. A rather general method to distinguish ordered from disordered phases in a quantum system is to study the reaction of the ground state to changes in boundary conditions. Therefore we investigate the low-energy states of the model using randomly twisted boundary conditions (RTBCs), following ideas introduced in Ref. [58]. Specifically, we study how changes by randomly twisting the boundary conditions affect the ground state. The twisting is performed by adding a complex phase to the hopping terms that cross a boundary in the horizontal or vertical direction. A hard-core boson crossing the boundary in the x direction then acquires a phase θ_x , for a vertical hop across

the boundary it picks up a phase θ_y , and it picks up the sum or difference $\theta_x \pm \theta_y$ in the case of a diagonal crossing.

For each realization, the phases (θ_x, θ_y) are drawn at random from the uniformly distributed interval $[0, 2\pi)$. For further reference we will use P to denote the set of realizations, where

$$\{\theta_x^{(p)}, \theta_y^{(p)}\} \in P, \quad \forall 0 \leq p \leq M, \quad (15)$$

and M represents the number of realizations. Depending on the particular boundary condition in one realization, the ground-state energy $E(\theta_x^{(p)}, \theta_y^{(p)})$ and the ground-state vector $|\Psi(\theta_x^{(p)}, \theta_y^{(p)})\rangle$ will vary, some realizations resulting in higher or lower ground-state energies. Therefore we define the *optimal twist* $(\theta_x^{gs}, \theta_y^{gs})$ to be that realization which results in the minimal ground-state energy

$$E(\theta_x^{gs}, \theta_y^{gs}) \equiv \min_{p \in P} \{E(\theta_x^{(p)}, \theta_y^{(p)})\}. \quad (16)$$

Accordingly, we will refer to $|\Psi(\theta_x^{gs}, \theta_y^{gs})\rangle$ as the *optimal ground-state vector* and to $E(\theta_x^{gs}, \theta_y^{gs})$ as the *optimal ground-state energy*. Then, we can normalize all energies $E(\theta_x^{(p)}, \theta_y^{(p)})$ with respect to the optimal ground-state energy and compute the relative difference

$$\epsilon_p = \frac{E(\theta_x^{(p)}, \theta_y^{(p)}) - E(\theta_x^{gs}, \theta_y^{gs})}{|E(\theta_x^{gs}, \theta_y^{gs})|} > 0, \quad (17)$$

as well as the overlap of each ground state with the optimal ground state

$$O_p = | \langle \Psi(\theta_x^{(p)}, \theta_y^{(p)}) | \Psi(\theta_x^{gs}, \theta_y^{gs}) \rangle |. \quad (18)$$

As the authors explain in Ref. [56], the distribution of O_p over ϵ_p for a set of ground-state vectors (all relevant parameters of the Hamiltonian remaining the same) depends strongly on whether a quantum phase is ordered or disordered. For an ordered phase, there exists a definitive boundary condition which accommodates the order intrinsic to the ground state of an infinite system, whereas all other boundary conditions prohibit it. An antiferromagnetic spin- $\frac{1}{2}$ chain in 1D is a simple example. Here, if the number of sites in the system is odd, the antiferromagnetic order is prevented if no twisted boundary conditions are in place. Including TBCs, $\theta = \pi$ is uniquely suited to minimize the ground-state energy as it accommodates the order of the system. As θ is altered from its optimal value of π , the ground-state energy increases. Therefore we expect only very few ground states of similar energy to the optimal ground-state energy for an ordered phase. In the disordered case, however, many different boundary conditions lead to very similar ground-state energies, including states that have very little overlap with the optimal ground state. To quantify this distribution, we define the set of configurations with energy comparable to the optimal ground state

$$Q = \{(\theta_x^{(p)}, \theta_y^{(p)}) \in P : \epsilon_p < \alpha\}, \quad (19)$$

where α is chosen to be sufficiently small, e.g., $\alpha = 0.01$. Subsequently, we define the fraction of low-energy configurations to be

$$N_c = \frac{|Q|}{M}. \quad (20)$$




FIG. 10. Results of the RTBC calculation for (a) $g = -5$, (b) $g = 0$, (c) $g = 0.73$, and (d) $g = 4.0$. As detailed in the text, $N_c \approx 1$ signals a disordered phase, where $N_c \approx 0$ is expected for an ordered state. Consequently, we identify phase I as a possible spin-liquid candidate regime, whereas regime II shows no distinct features.

Continuing the argument from before, N_c is typically small for ordered phases and close to unity for disordered ones. This characterization is somewhat dependent on the particular choice of α , which also depends on the system size L . For our case of $L = 24$ (we use the shape 24C shown in Fig. 2) the choice of $\alpha = 0.01$ is reasonable (see Ref. [56]).

In Fig. 10 we show the results of the RTBC calculations for shape 24C, having performed $M = 200$ realizations of twisting angles for each value of g that we consider. In Fig. 10(d), not all points are visible since we limit the ϵ_p axis to 0.05. Judging by the values of N_c for each g , we find a clearly disordered regime for $g = -5$ (regime I in Fig. 5), with a strongly ordered phase at $g = 4$ (120° order). The RTBC results at value $g = 0.73$, i.e., in phase II, are in between the clearly disordered situation shown in Fig. 10(a) and the BEC phase, shown in Fig. 10(b), and a clear identification as an ordered or disordered phase is not easy. We will show, however, in Sec. III C that the situation becomes much clearer if the competing effects from the repulsive density-density interaction are switched off.

C. Hamiltonian without density-density interaction

The character of phase II is somewhat masked by the simultaneous presence of a density-density interaction, which drives the system into trivial ordered states. Therefore we consider a modified version of Hamiltonian (1), where we artificially switch off the density-density interaction:

$$\hat{H} = -J \sum_{\langle i,j \rangle} \hat{b}_j^\dagger \hat{b}_i - 2gJ \sum_{\langle\langle i,j \rangle\rangle} \hat{b}_j^\dagger \hat{b}_i e^{\pm \frac{2\pi i}{3}} (1 - \hat{n}_{ij}). \quad (21)$$

We first verify that the emergence of phases I and II is not due to density-density interactions. This can be seen from Fig. 11, where we have plotted the ground-state fidelity metric for Hamiltonian (21). In addition to the two peaks at small positive g we find that the slow crossover to regime I at large negative g is shifted to even larger absolute values. Without the density-density interaction, the system is less likely to show long-range diagonal order. For this reason, we again consider the sensitivity to changes in boundary conditions (RTBCs). The corresponding results are shown in Fig. 12. We observe




FIG. 11. Ground-state fidelity metric f as a function of the parameter g for the Hamiltonian (21). The peaks of f agree well for the different shapes and indicate potential phase transitions. The inset shows regime II in detail. We find the same qualitative behavior as for Hamiltonian (1), but the transition points are shifted.

that the intermediate phase II is now characterized by a large value of N_c , clearly signaling a disordered ground state, while N_c stays roughly the same for the other phases.

IV. NATURE OF THE LIQUID PHASE

Having established the liquidlike behavior of phase II, we will investigate in the following the possible nature of this phase.

A. Many-body Chern number

The complex NNN hopping in the Hamiltonian explicitly breaks time-reversal symmetry. We now show that as a consequence of this and of the nonlinear character of the NNN hopping, phase II is topological, characterized by a nonvanishing Chern number. The many-body Chern number in a two-dimensional lattice model on a torus can conveniently be obtained from the many-body ground-state wave function $|\Psi(\theta)\rangle = |\Psi(\theta^x, \theta^y)\rangle$ with twisted boundary conditions in the




FIG. 12. Results of the RTBC calculation for (a) $g = -15$, (b) $g = 0$, (c) $g = 0.42$, and (d) $g = 4.0$ for Hamiltonian (21). As detailed in the text, $N_c \approx 1$ signals a disordered phase, where $N_c \approx 0$ is expected for an ordered state. We observe that regime II now shows strong disorder whereas for the full Hamiltonian we saw a distribution of ground states closer to the BEC case.




FIG. 13. Many-body Zak phase winding for $g = 0.73$ and the trivial cases for $g = 0$ (BEC) and $g = 4.0$ (120°), calculated for the full Hamiltonian (1).

x and y directions, respectively:

$$C = \frac{i}{2\pi} \int_0^{2\pi} d\theta_x \int_0^{2\pi} d\theta_y (\langle \partial_{\theta_x} \Psi(\theta) | \partial_{\theta_y} \Psi(\theta) \rangle - \text{c.c.}). \quad (22)$$

For the numerical calculation we use a set of discrete twisting angles $\{\theta_i^x, \theta_j^y\}$, where

$$\theta_i^\alpha = \frac{2\pi}{D} i, \quad (23)$$

with $\alpha = x, y$ and D being the number of intervals. We then calculate the ground-state wave function $|\Psi(\theta_i^x, \theta_j^y)\rangle$ for each $\{\theta_i^x, \theta_j^y\}$ and calculate the many-body Zak phase using

$$\begin{aligned} \phi^{\text{MB}}(\theta_j^y) = & \text{Im} \ln [\langle \Psi(\theta_1^x, \theta_j^y) | \Psi(\theta_2^x, \theta_j^y) \\ & \times \cdots \langle \Psi(\theta_D^x, \theta_j^y) | \Psi(\theta_1^x, \theta_j^y) \rangle]. \end{aligned} \quad (24)$$

Here, we take the loop product in the x twisting angle only. The Chern number can then be calculated as the winding of the many-body Zak phase

$$C = \frac{1}{2\pi} \int_0^{2\pi} d\theta^y \frac{\partial \phi^{\text{MB}}}{\partial \theta^y}. \quad (25)$$

In doing so, we obtain for $g = 0.73$ a Chern number of $C = 1$ to within numerical precision (see Fig. 13). We checked that the Chern number did not change when adding a potential disorder of $\pm 0.1J$. We also checked an additional shape (shape 20A) and again obtain a Chern number of $C = 1$. For comparison, the calculation of the Chern number in the 120° phase, i.e., for $g = 4$, yields $C = 0$ again to within numerical precision.

We did not calculate a Chern number for the disordered and gapless regime I, as varying the twisting angles mixes the ground state with excited states.

B. Spin gap and collective gap

In the study of disordered spin states, the distinction between gapless and gapped spin liquids is important. Therefore we now investigate the spin gap and the collective gap.

In a hard-core boson representation the spin gap corresponds to the change in the energy per particle when adding or subtracting a boson. Thus we calculate the chemical potential as the discrete first derivative of the many-body energy with




FIG. 14. (a) Chemical potential for the full Hamiltonian (1) on shape 24C in regimes I and II, plotted over the density N/L , where N represents the number of particles in the system. (b) and (c) Energy of the lowest excited state, normalized with respect to the ground state, for cluster shape 24C and $N/L = \frac{1}{2}$ for regimes I and II. In phase II, one recognizes a nondegenerate ground state. In phase I the energy gap is an order of magnitude smaller than in phase II. The absolute energies per particle which we obtain at $g = -5$ are $E \approx -5.493 |gJ|$, and at $g = 0.73$ we find $E \approx -0.763 |gJ|$.

regard to the particle number N . The result for the disordered regimes I and II can be seen in Fig. 14(a). For the small systems that we are able to analyze numerically with exact diagonalization, the clear identification of a spin gap is masked by finite-size effects. Nevertheless, while in phase I the spin gap clearly vanishes [see Fig. 14(a)], the curve for phase II is indicative of a finite spin gap.

Secondly, we investigate the collective gap, which is shown in Fig. 14(b) for the parameter regimes of the BEC and phase I and in Fig. 14(c) for phase II. We plot the energy gap to the first excited state in the mean-field Hamiltonian (red symbols), the full Hamiltonian (blue symbols), and the Hamiltonian without density-density interaction terms (green symbols) for cluster shape 24C. We have also calculated the collective gap for a few values of g in clusters up to size 30 but could not make a reliable finite-size scaling. Phase I is clearly gapless, while for phase II we observe that the excitation gap is increased by the nonlinear hopping as compared with the mean-field case, and, as expected, the density-density interaction leads to an additional enhancement. This is indicative of a gapped phase II and thus a nondegenerate ground state on a torus.

The absence of degeneracy in the topological nontrivial regime would point to a symmetry-protected topological (SPT) phase. Our model has a $U(1)$ symmetry associated with particle-number conservation. There is a full classification of SPT phases [25], and for bosons in $d = 2$ spatial dimensions with $U(1)$ symmetry, different phases characterized by a \mathbf{Z} -quantized topological invariant exist, corresponding to the $\mathcal{H}^{1+d}[U(1), U(1)]$ cohomology group [25]. We note, however, that our finding of an odd-valued Chern number is different from the Chern numbers $C = \pm 2$ of the bosonic integer quantum Hall (BIQH) effect found, e.g., in Ref. [38] for bosons on a honeycomb lattice with NN and (different) density-dependent NNN hopping at unit filling and in Ref. [37] or Ref. [39] for bosons with internal degrees of freedom. The odd value of C is also different from what would be expected from the classification of interacting integer topological phases put forward in Ref. [24].

We are not able to perform a proper finite-size scaling using ED simulations, and for the unambiguous verification

of the gapfulness of phase II, more sophisticated methods are needed. Corresponding investigations using DMRG and a tensor network approach are underway and will be reported elsewhere [35]. In particular, for a Dirac QSL it is difficult to verify the gaplessness of the system, but the dependence of the collective gap on twisting angles is a signature of such a gapless spin liquid [59]. In Fig. 15 we have therefore analyzed the size of the gap for different twists of the boundary conditions. While the gap remains always finite, it shows a strong dependence.

C. Fermion Hamiltonian and Chern-Simons gauge field

In the following we argue that the origin of the topological phase II can be understood from a representation of the model in terms of fermions. A mapping from hard-core bosons to spinless fermions can be achieved in one dimension by a Jordan-Wigner transformation [60]. In two dimensions, this is accomplished via a Chern-Simons (CS) transformation, whose lattice version [61] reads

$$\hat{b}_i = \hat{c}_i e^{i \sum_{j \neq i} \arg(z_i - z_j)} \hat{n}_j, \tag{26}$$




FIG. 15. Collective gap as a function of twisted boundary angles θ_x and θ_y in phase II for shape 24C, normalized to the gap at zero twist. Left: For Hamiltonian (21), i.e., without density-density interaction, and $g = 0.42$. Right: For the full Hamiltonian (1) and $g = 0.73$.

where \hat{b}_j are hard-core boson operators and \hat{c}_j are fermion operators. Here, $z_j = x_j + iy_j$ are the complex positions in the 2D lattice. (Note that we use a different sign convention from that in Ref. [61].) When applying the CS transformation to our Hamiltonian, we find

$$\begin{aligned} \hat{H} = & -J \sum_{\langle i,j \rangle} \hat{c}_j^\dagger \hat{c}_i e^{i\hat{B}_{ji}} - 2gJ \sum_{\langle\langle i,j \rangle\rangle} \hat{c}_j^\dagger \hat{c}_i e^{\pm \frac{2\pi i}{3}} e^{i\hat{B}_{ji}} (1 - \hat{n}_{ij}) \\ & + 2gJ \sum_{\langle i,j \rangle} \hat{n}_i \hat{n}_j, \end{aligned} \quad (27)$$

where a Chern-Simons gauge field

$$\hat{B}_{ji} = \sum_{l \neq j,i} [\arg(z_i - z_l) - \arg(z_j - z_l)] \hat{n}_l \quad (28)$$

appears. It is instructive to decompose this field into a mean-field part and a fluctuation part $\hat{B}_{ji} = \langle \hat{B}_{ji} \rangle + \delta \hat{B}_{ji}$, where $\delta \hat{B}_{ji} = \sum_{l \neq j,i} \delta \hat{B}_{ji}^{(l)}$ and

$$\delta \hat{B}_{ji}^{(l)} = [\arg(z_i - z_l) - \arg(z_j - z_l)] (\hat{n}_l - \langle \hat{n}_l \rangle). \quad (29)$$

The mean-field term can easily be evaluated for an infinite hexagonal lattice at half filling, where $\langle \hat{n}_j \rangle = 0.5$. One finds that $\langle \hat{B}_{ji} \rangle$ is to good approximation a multiple of 2π for the NN hopping terms and may be disregarded, while for the NNN hoppings it just compensates the terms $\pm 2\pi/3$ to within a few percent. Thus the system can approximately be described by a Haldane model for fermions in the topological trivial regime of real-valued NNN hoppings which interact with a fluctuation Chern-Simons field and have a density-dependent NNN hopping.

$$\begin{aligned} \hat{H} \approx & -J \sum_{\langle i,j \rangle} \hat{c}_j^\dagger \hat{c}_i e^{i\delta \hat{B}_{ji}} - 2gJ \sum_{\langle\langle i,j \rangle\rangle} \hat{c}_j^\dagger \hat{c}_i e^{i\delta \hat{B}_{ji}} (1 - \hat{n}_{ij}) \\ & + 2gJ \sum_{\langle i,j \rangle} \hat{n}_i \hat{n}_j. \end{aligned} \quad (30)$$

We have seen in Sec. III A that within a mean-field approximation of the projector $(1 - \hat{n}_{ij}) \rightarrow (1 - \bar{n})$, the topological phase II disappears. We thus conclude that here the fluctuation CS field $\delta \hat{B}_{ji}$ can most likely be neglected. In the full model, however, the projector $(1 - \hat{n}_{ij})$ generates an additional mean-field contribution resulting from the site in between the next-nearest neighbors i and j . Denoting this site in here as $l = 0$, we find

$$\begin{aligned} & \exp \left(i \sum_{l \neq j,i} \delta \hat{B}_{ji}^{(l)} \right) (1 - \hat{n}_0) \\ & = \exp \left(i \sum_{l \neq j,i,0} \delta \hat{B}_{ji}^{(l)} \right) e^{\pm \frac{\pi i}{3}} (1 - \hat{n}_0) \approx e^{\pm \frac{\pi i}{3}} (1 - \hat{n}_0). \end{aligned} \quad (31)$$

Assuming that all other contributions to the fluctuation CS field $\delta \hat{B}_{ji}^{(l)}$ are small and can be ignored, we recognize that the nonlinearity of the NNN hopping effectively generates a nonvanishing flux for the fermions, which therefore enter a topologically nontrivial phase of the Haldane model with Chern number $C = 1$. Thus we identify as the origin of the




FIG. 16. Comparison of ground-state fidelity as a function of g for cluster shape 24C with (red, dashed curve) and without (blue, solid curve) direct NNN hopping, resulting from expression (32).

topologically nontrivial phase II the additional Chern-Simons field created by the nonlinearity in the NNN hopping.

V. EXPERIMENTAL CONSIDERATIONS: EFFECTS OF LONGER-RANGE INTERACTIONS

One of the main motivations of our work is to show that Rydberg excitations in an array of trapped atoms are a suitable platform to observe spin liquids. We have modeled the system with a simplified Hamiltonian (1), and thus some comments are in order about the limitations of this model. The microscopic origin of the direct hopping of spin excitations is the resonant exchange of a microwave photon giving rise to an $\sim 1/r^3$ dipole-dipole coupling. In the Hamiltonian (1) we have only taken into account direct exchange couplings between nearest neighbors, neglecting the coupling to next-nearest neighbors. These processes would lead to an additional NNN hopping contribution to the Hamiltonian

$$\hat{H}_{\text{LR}} = -\frac{J}{(\sqrt{3})^3} \sum_{\langle\langle i,j \rangle\rangle} \hat{b}_j^\dagger \hat{b}_i, \quad (32)$$

which does not affect phase I and the 120° phase for $|g| \rightarrow \infty$. It is, however, a sizable modification of the NNN hopping in the regime of $g \lesssim 1$. We have numerically checked that the inclusion of (32) does not compromise the emergence of the nontrivial regime II and only leads to a quantitative shift of the critical values for g . This is illustrated in Fig. 16, where we have compared the ground state fidelity with and without \hat{H}_{LR} .

VI. SUMMARY AND CONCLUSION

Despite numerous experimental indications, the realization and verification of a quantum spin liquid phase in solid-state systems remain a major challenge. In this paper we have proposed a model system accessible in cold-gas experiments with Rydberg atoms where such a state could be realized and studied in extended parameter regimes. Motivated by the recent experimental observation of nonlinear, complex hopping processes of Rydberg spin excitations in two-dimensional arrays of trapped atoms, we analyzed the many-body ground state of these excitations in a honeycomb lattice at half filling using exact diagonalization simulations. The density-dependent complex hopping as well as a nearest-

neighbor density-density interaction arise from second-order processes of excitations between two Rydberg levels, whose strength can be controlled by tuning the energy of a third, off-resonant Rydberg state. If the nonlinear hopping is treated in mean-field approximation, the model is equivalent to the Haldane model in the topologically nontrivial phase with additional nearest-neighbor interactions. Since the elementary constituents are here spin- $\frac{1}{2}$ particles or, equivalently, hard-core bosons rather than fermions, the mean-field model has, however, no topological ground state. It instead possesses only two trivial phases, a condensate (BEC) with a preferred occupation of modes with wave vectors close to zero, as well as a 120° phase with spiral spin order and a remaining $SO(2)$ rotational symmetry. A phase transition between the two phases occurs when the direct hopping is similar in amplitude to the second-order, next-nearest-neighbor hopping. In the full model an additional phase, denoted as phase II, emerges close to the mean-field critical point, and there are indications of a transition or a crossover into another phase for very large second-order hoppings of opposite sign. The latter regime is, however, outside the range of validity of the effective many-body Hamiltonian for the Rydberg system. We verified the absence of simple spin or dimer order in the new phases, and considering randomly twisted boundary conditions, we found strong evidence that both phases are disordered. This becomes particularly evident if the density-density interaction, which competes with the hopping processes and drives the system towards density order, is switched off. Since the complex, nonlinear hopping breaks both time-reversal and chiral symmetry, the spins can have a nonvanishing spin chirality, which attains a very large value in both new phases. We calculated the many-body Chern number and found a value $C = 1$ to within numerical precision in phase II, which was also shown to be robust against potential disorder, and $C = 0$ in the BEC and 120° phases. Furthermore, we calculated the spin and collective gaps using ED simulations on finite lattices with periodic boundary conditions. Since for the system sizes that can be reached with exact diagonalization it is not possible to make reliable extrapolations to the thermodynamic limit, we cannot draw definite conclusions here. While the ED

results clearly point to a gapless phase I, we found indications for a finite spin gap and a finite collective gap in phase II, which was shown to originate from the nonlinear hopping rather than from the density-density interaction. This would point to a symmetry-protected topological phase protected by the $U(1)$ symmetry. Considering a mapping of the hard-core boson Hamiltonian to spinless fermions coupled to a Chern-Simons field, we showed that the topologically nontrivial phase is caused by the density dependence of the NNN hopping. The nonlinearity of this hopping generates an additional Chern-Simons flux for the fermion model, which becomes topologically nontrivial due to this. The odd value $C = 1$ of the Chern number is, however, in contrast to several bosonic integer quantum Hall phases found in Refs. [37–39] and predicted from the general classification scheme of Ref. [24]. Calculating the collective gap in phase II for different twisted boundary conditions, we found a nonzero, but strongly varying value, which rather points to a gapless Dirac QSL, which is even more pronounced if the density-density interaction is switched off. Thus while there is strong evidence that phase II is a topological quantum spin liquid at half filling, its true nature remains unclear and requires further investigation.

Note added. Recently, we became aware of a publication [62] predicting a fractional quantum Hall phase for a similar system with average particle density $\frac{1}{4}$ induced by density-density interactions in engineered flat Chern bands.

ACKNOWLEDGMENTS

We thank Marcello Dalmonte, Poetri Tarabunga, Titas Chanda, and Giuliano Giudici for very fruitful and inspiring discussions. We furthermore thank Frank Pollmann and Tigran Sedrakyan for useful comments. The authors gratefully acknowledge financial support from the DFG through SFB TR 185, Project No. 277625399.

M.F. conceived and supervised the project; S.O. performed the analytic calculations and the numerical simulations; S.O. and M.K.-E. worked on the initial numerical implementation; all authors analyzed the data; S.O. and M.F. worked on the final manuscript with input from M.K.-E.

-
- [1] H. Weimer, M. Müller, I. Lesanovsky, P. Zoller, and H. P. Büchler, A Rydberg quantum simulator, *Nat. Phys.* **6**, 382 (2010).
 - [2] P. Schauß, M. Cheneau, M. Endres, T. Fukuhara, S. Hild, A. Omran, T. Pohl, C. Gross, S. Kuhr, and I. Bloch, Observation of spatially ordered structures in a two-dimensional Rydberg gas, *Nature (London)* **491**, 87 (2012).
 - [3] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, Probing many-body dynamics on a 51-atom quantum simulator, *Nature (London)* **551**, 579 (2017).
 - [4] A. Browaeys and T. Lahaye, Many-body physics with individually controlled Rydberg atoms, *Nat. Phys.* **16**, 132 (2020).
 - [5] F. M. Surace, P. P. Mazza, G. Giudici, A. Lerose, A. Gambassi, and M. Dalmonte, Lattice Gauge Theories and String Dynamics in Rydberg Atom Quantum Simulators, *Phys. Rev. X* **10**, 021041 (2020).
 - [6] P. Scholl, M. Schuler, H. J. Williams, A. A. Eberharter, D. Barredo, K.-N. Schymik, V. Lienhard, L.-P. Henry, T. C. Lang, T. Lahaye, A. M. Läuchli, and A. Browaeys, Quantum simulation of 2D antiferromagnets with hundreds of Rydberg atoms, *Nature (London)* **595**, 233 (2021).
 - [7] D. Jaksch, J. I. Cirac, P. Zoller, S. L. Rolston, R. Côté, and M. D. Lukin, Fast Quantum Gates for Neutral Atoms, *Phys. Rev. Lett.* **85**, 2208 (2000).
 - [8] M. D. Lukin, M. Fleischhauer, R. Cote, L. M. Duan, D. Jaksch, J. I. Cirac, and P. Zoller, Dipole Blockade and Quantum Information Processing in Mesoscopic Atomic Ensembles, *Phys. Rev. Lett.* **87**, 037901 (2001).

- [9] A. Gaëtan, Y. Miroshnychenko, T. Wilk, A. Chotia, M. Viteau, D. Comparat, P. Pillet, A. Browaeys, and P. Grangier, Observation of collective excitation of two individual atoms in the Rydberg blockade regime, *Nat. Phys.* **5**, 115 (2009).
- [10] E. Urban, T. A. Johnson, T. Henage, L. Isenhower, D. Yavuz, T. Walker, and M. Saffman, Observation of Rydberg blockade between two atoms, *Nat. Phys.* **5**, 110 (2009).
- [11] M. Saffman, T. G. Walker, and K. Mølmer, Quantum information with Rydberg atoms, *Rev. Mod. Phys.* **82**, 2313 (2010).
- [12] S. de Léséleuc, V. Lienhard, P. Scholl, D. Barredo, S. Weber, N. Lang, H. P. Büchler, T. Lahaye, and A. Browaeys, Observation of a symmetry-protected topological phase of interacting bosons with Rydberg atoms, *Science* **365**, 775 (2019).
- [13] G. Semeghini, H. Levine, A. Keesling, S. Ebadi, T. T. Wang, D. Bluvstein, R. Verresen, H. Pichler, M. Kalinowski, R. Samajdar, A. Omran, S. Sachdev, A. Vishwanath, M. Greiner, V. Vuletić, and M. D. Lukin, Probing topological spin liquids on a programmable quantum simulator, *Science* **374**, 1242 (2021).
- [14] P. W. Anderson, Resonating valence bonds: A new kind of insulator? *Mater. Res. Bull.* **8**, 153 (1973).
- [15] P. W. Anderson, The resonating valence bond state in La_2CuO_4 and superconductivity, *Science* **235**, 1196 (1987).
- [16] F. Mila, Quantum spin liquids, *Eur. J. Phys.* **21**, 499 (2000).
- [17] P. A. Lee, An end to the drought of quantum spin liquids, *Science* **321**, 1306 (2008).
- [18] L. Savary and L. Balents, Quantum spin liquids: a review, *Rep. Prog. Phys.* **80**, 016502 (2017).
- [19] C. Broholm, R. Cava, S. Kivelson, D. Nocera, M. Norman, and T. Senthil, Quantum spin liquids, *Science* **367**, eaay0668 (2020).
- [20] Y. Shen, Y.-D. Li, H. Wo, Y. Li, S. Shen, B. Pan, Q. Wang, H. C. Walker, P. Steffens, M. Boehm, Y. Hao, D. L. Quintero-Castro, L. W. Harriger, M. D. Frontzek, L. Hao, S. Meng, Q. Zhang, G. Chen, and J. Zhao, Evidence for a spinon Fermi surface in a triangular-lattice quantum-spin-liquid candidate, *Nature (London)* **540**, 559 (2016).
- [21] R. Coldea, D. A. Tennant, A. M. Tsvelik, and Z. Tylczynski, Experimental Realization of a 2D Fractional Quantum Spin Liquid, *Phys. Rev. Lett.* **86**, 1335 (2001).
- [22] J. Liu, L. Yuan, X. Li, B. Li, K. Zhao, H. Liao, and Y. Li, Gapless spin liquid behavior in a kagome Heisenberg antiferromagnet with randomly distributed hexagons of alternate bonds, *Phys. Rev. B* **105**, 024418 (2022).
- [23] M. Gohlke, G. Wachtel, Y. Yamaji, F. Pollmann, and Y. B. Kim, Quantum spin liquid signatures in Kitaev-like frustrated magnets, *Phys. Rev. B* **97**, 075126 (2018).
- [24] Y.-M. Lu and A. Vishwanath, Theory and classification of interacting integer topological phases in two dimensions: A Chern-Simons approach, *Phys. Rev. B* **86**, 125119 (2012); **89**, 199903(E) (2014).
- [25] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Symmetry protected topological orders and the group cohomology of their symmetry group, *Phys. Rev. B* **87**, 155114 (2013).
- [26] X. Chen, Z.-C. Gu, and X.-G. Wen, Local unitary transformation, long-range quantum entanglement, wave function renormalization, and topological order, *Phys. Rev. B* **82**, 155138 (2010).
- [27] X.-G. Wen, A theory of 2 + 1D bosonic topological orders, *Natl. Sci. Rev.* **3**, 68 (2016).
- [28] J. Knolle and R. Moessner, A field guide to spin liquids, *Annu. Rev. Condens. Matter Phys.* **10**, 451 (2019).
- [29] J. Wen, S.-L. Yu, S. Li, W. Yu, and J.-X. Li, Experimental identification of quantum spin liquids, *npj Quantum Mater.* **4**, 12 (2019).
- [30] M. R. Norman, Colloquium: Herbertsmithite and the search for the quantum spin liquid, *Rev. Mod. Phys.* **88**, 041002 (2016).
- [31] R. Verresen, M. D. Lukin, and A. Vishwanath, Prediction of Toric Code Topological Order from Rydberg Blockade, *Phys. Rev. X* **11**, 031005 (2021).
- [32] R. Samajdar, W. W. Ho, H. Pichler, M. D. Lukin, and S. Sachdev, Quantum phases of Rydberg atoms on a kagome lattice, *Proc. Natl. Acad. Sci. USA* **118**, e2015785118 (2021).
- [33] V. Lienhard, P. Scholl, S. Weber, D. Barredo, S. de Léséleuc, R. Bai, N. Lang, M. Fleischhauer, H. P. Büchler, T. Lahaye, and A. Browaeys, Realization of a Density-Dependent Peierls Phase in a Synthetic, Spin-Orbit Coupled Rydberg System, *Phys. Rev. X* **10**, 021031 (2020).
- [34] F. Görg, K. Sandholzer, J. Minguzzi, R. Desbuquois, M. Messer, and T. Esslinger, Realization of density-dependent Peierls phases to engineer quantized gauge fields coupled to ultracold matter, *Nat. Phys.* **15**, 1161 (2019).
- [35] M. Dalmonte (private communication).
- [36] T. Senthil and M. Levin, Integer Quantum Hall Effect for Bosons, *Phys. Rev. Lett.* **110**, 046801 (2013).
- [37] A. Sterdyniak, N. R. Cooper, and N. Regnault, Bosonic Integer Quantum Hall Effect in Optical Flux Lattices, *Phys. Rev. Lett.* **115**, 116802 (2015).
- [38] Y.-C. He, S. Bhattacharjee, R. Moessner, and F. Pollmann, Bosonic Integer Quantum Hall Effect in an Interacting Lattice Model, *Phys. Rev. Lett.* **115**, 116803 (2015).
- [39] W. Liu, Z. Dong, Z. Dong, C. Liu, W. Yan, and Y. Chen, Bosonic integer quantum Hall state without Landau levels on a square lattice, *Phys. Rev. B* **99**, 085305 (2019).
- [40] C. N. Varney, K. Sun, V. Galitski, and M. Rigol, Kaleidoscope of Exotic Quantum Phases in a Frustrated XY Model, *Phys. Rev. Lett.* **107**, 077201 (2011).
- [41] C. N. Varney, K. Sun, V. Galitski, and M. Rigol, Quantum phases of hard-core bosons in a frustrated honeycomb lattice, *New J. Phys.* **14**, 115028 (2012).
- [42] Z. Zhu, D. A. Huse, and S. R. White, Unexpected z -Direction Ising Antiferromagnetic Order in a Frustrated Spin-1/2 $J_1 - J_2XY$ Model on the Honeycomb Lattice, *Phys. Rev. Lett.* **111**, 257201 (2013).
- [43] M. Endres, H. Bernien, A. Keesling, H. Levine, E. R. Anschuetz, A. Krajenbrink, C. Senko, V. Vuletic, M. Greiner, and M. D. Lukin, Atom-by-atom assembly of defect-free one-dimensional cold atom arrays, *Science* **354**, 1024 (2016).
- [44] D. Barredo, S. de Léséleuc, V. Lienhard, T. Lahaye, and A. Browaeys, An atom-by-atom assembler of defect-free arbitrary two-dimensional atomic arrays, *Science* **354**, 1021 (2016).
- [45] D. Barredo, V. Lienhard, S. de Léséleuc, T. Lahaye, and A. Browaeys, Synthetic three-dimensional atomic structures assembled atom by atom, *Nature (London)* **561**, 79 (2018).
- [46] S. Weber, S. De Léséleuc, V. Lienhard, D. Barredo, T. Lahaye, A. Browaeys, and H. P. Büchler, Topologically protected edge states in small Rydberg systems, *Quantum Sci. Technol.* **3**, 044001 (2018).

- [47] S. Ohler, M. Kiefer-Emmanouilidis, A. Browaeys, H. P. Büchler, and M. Fleischhauer, Self-generated quantum gauge fields in arrays of Rydberg atoms, *New J. Phys.* **24**, 023017 (2022).
- [48] P. Virtanen *et al.* (SciPy 1.0 Contributors), SciPy 1.0: fundamental algorithms for scientific computing in Python, *Nat. Methods* **17**, 261 (2020); **17**, 352(E) (2020).
- [49] P. Zanardi and N. Paunković, Ground state overlap and quantum phase transitions, *Phys. Rev. E* **74**, 031123 (2006).
- [50] L. Campos Venuti, M. Cozzini, P. Buonsante, F. Massel, N. Bray-Ali, and P. Zanardi, Fidelity approach to the Hubbard model, *Phys. Rev. B* **78**, 115410 (2008).
- [51] M.-F. Yang, Ground-state fidelity in one-dimensional gapless models, *Phys. Rev. B* **76**, 180403(R) (2007).
- [52] F. D. M. Haldane, Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the “Parity Anomaly”, *Phys. Rev. Lett.* **61**, 2015 (1988).
- [53] C. N. Varney, K. Sun, M. Rigol, and V. Galitski, Interaction effects and quantum phase transitions in topological insulators, *Phys. Rev. B* **82**, 115125 (2010).
- [54] E. Fradkin, Jordan-Wigner Transformation for Quantum-Spin Systems in Two Dimensions and Fractional Statistics, *Phys. Rev. Lett.* **63**, 322 (1989).
- [55] C. Hickey, L. Cincio, Z. Papić, and A. Paramekanti, Haldane-Hubbard Mott Insulator: From Tetrahedral Spin Crystal to Chiral Spin Liquid, *Phys. Rev. Lett.* **116**, 137202 (2016).
- [56] D. Castells-Graells, A. Yuste, and A. Sanpera, Frustrated quantum spin systems in small triangular lattices studied with a numerical method, *Phys. Rev. B* **100**, 155119 (2019).
- [57] Z. Meng, T. Lang, S. Wessel, F. Assaad, and A. Muramatsu, Quantum spin liquid emerging in two-dimensional correlated Dirac fermions, *Nature (London)* **464**, 847 (2010).
- [58] M. Thesberg and E. S. Sørensen, Exact diagonalization study of the anisotropic triangular lattice Heisenberg model using twisted boundary conditions, *Phys. Rev. B* **90**, 115117 (2014).
- [59] Y.-C. He, M. P. Zaletel, M. Oshikawa, and F. Pollmann, Signatures of Dirac Cones in a DMRG Study of the Kagome Heisenberg Model, *Phys. Rev. X* **7**, 031020 (2017).
- [60] P. Jordan and E. Wigner, Über das paulische Äquivalenzverbot, *Z. Phys.* **47**, 631 (1928).
- [61] T. A. Sedrakyan, L. I. Glazman, and A. Kamenev, Spontaneous Formation of a Nonuniform Chiral Spin Liquid in a Moat-Band Lattice, *Phys. Rev. Lett.* **114**, 037203 (2015).
- [62] S. Weber, R. Bai, N. Makki, J. Mögerle, T. Lahaye, A. Browaeys, M. Daghofer, N. Lang, and H. P. Büchler, Experimentally Accessible Scheme for a Fractional Chern Insulator in Rydberg Atoms, *PRX Quantum* **3**, 030302 (2022).