

Home Search Collections Journals About Contact us My IOPscience

Superfluid phases of spin-1 bosons in a cubic optical lattice

This content has been downloaded from IOPscience. Please scroll down to see the full text. 2013 Laser Phys. Lett. 10 115501

(http://iopscience.iop.org/1612-202X/10/11/115501)

View the table of contents for this issue, or go to the journal homepage for more

Download details: This content was downloaded by: pelster IP Address: 130.133.31.70 This content was downloaded on 20/10/2013 at 08:21

Please note that terms and conditions apply.

Laser Phys. Lett. 10 (2013) 115501 (6pp)

LETTER

Superfluid phases of spin-1 bosons in a cubic optical lattice

Mohamed Mobarak¹ and Axel Pelster^{2,3}

¹ Insitute für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, D-14195 Berlin, Germany

² Fachbereich Physik und Forschungszentrum OPTIMAS, Technische Universität Kaiserslautern,

D-67663 Kaiserslautern, Germany

³ Hanse-Wissenschaftskolleg, Lehmkuhlenbusch 4, D-27733 Delmenhorst, Germany

E-mail: axel.pelster@physik.uni-kl.de

Received 4 August 2013 Accepted for publication 4 September 2013 Published 16 October 2013 Online at stacks.iop.org/LPL/10/115501

Abstract

We analyze theoretically the emergence of different superfluid phases of spin-1 bosons in a three-dimensional cubic optical lattice by generalizing the recently developed Ginzburg–Landau theory for the Bose–Hubbard model to a spinor Bose gas. In particular at zero temperature, our theory distinguishes within its validity range between various superfluid phases for an anti-ferromagnetic interaction with an external magnetic field. In addition, we determine that the superfluid–Mott insulator phase transition is of second order and that the transitions between the respective superfluid phases with anti-ferromagnetic interaction can be both of first and second order.

(Some figures may appear in colour only in the online journal)

1. Introduction

In recent years optical lattices have become a major research topic within the realm of ultracold quantum gases, as they offer the perspective to simulate condensed matter physics under well-controlled conditions [1, 2]. Most prominently, the quantum phase transition between a superfluid (SF) and a Mott insulating (MI) phase of a spinless Bose gas loaded in a periodic optical potential has been experimentally observed by increasing the lattice depth [3, 4]. All properties of this quantum phase transition are captured by the underlying Bose-Hubbard Hamiltonian [5-7] for which different analytical solution methods have been worked out [8-14] and high-precision Monte Carlo studies have been performed [15, 16]. Furthermore, extensions of the Bose-Hubbard model have been investigated, which cover, for instance, superlattices [17], Bose–Fermi mixtures [18-21], quantum simulations like entanglement of atoms or quantum teleportation [22] disorder and [2, 23-25].

Preparing experimentally a spin-1 Bose-Einstein condensation (BEC) of ²³Na or ⁸⁷Rb atoms in an optical trap, the atomic spin degrees of freedom are not frozen due to the electric dipole force between atoms and the electric field of a laser beam [26, 27]. This experimental realization of an optically trapped BEC opened a new window to study also various phenomena of spinor Bose gases loaded in an optical lattice. For instance, they offer the possibility of studying strongly correlated states; for example the coherent collisional spin dynamics in an optical lattice was measured in [28] and the ⁸⁷Rb scattering lengths for F = 1 and 2 were determined in [29]. In particular, combining the spin degree of freedom with various types of interactions and with different lattice geometries offers the prospect to realize a plethora of superfluid phases with magnetic properties. A first tentative step in this direction was the loading of ⁸⁷Rb in a frustrated triangular lattice [30]. Despite these initial promising investigations, spinor Bose gases in an optical lattice seem experimentally to be so challenging that no further detailed experiments have been performed.

Theoretical progress in the study of spinor Bose gases in an optical lattice was made by [31, 32]. In the case of the anti-ferromagnetic interaction of ²³Na, the location of the SF-MI transition and several properties for SF and MI phases for spin-1 bosons were determined without external magnetic field at zero temperature. In particular, they found that the superfluid transition occurs into a polar spin-0 state [32] and the SF phase represents a polar state with zero spin expectation value. On the other hand, the effect of a non-vanishing external magnetic field upon the SF-MI transition was determined within a mean-field approximation in [33, 34]. In addition, it was also shown in [33, 34] that the superfluid transition occurs from the Mott insulating phase into either a polar spin-1 or spin-(-1) state, but investigation was not undertaken regarding which other phases might emerge deep in the superfluid phase.

In this letter we study the effect of the magnetic field on the emergence of superfluid phases of spin-1 bosons. We show that new superfluid phases can emerge due to the interplay of the anti-ferromagnetic interaction of spin-1 bosons and an external magnetic field in a three-dimensional cubic optical lattice at zero temperature. To this end, we extend the Ginzburg–Landau theory developed in [10, 11] from the spin-0 to the spin-1 Bose–Hubbard model. In the grand-canonical ensemble the underlying Hamiltonian can be decomposed according to $\hat{H}_{\rm BH} = \hat{H}^{(0)} + \hat{H}^{(1)}$ [31, 32], where the local part $\hat{H}^{(0)} = \sum_i \hat{H}_i^{(0)}$ reads

$$\hat{H}_{i}^{(0)} = \frac{U_{0}}{2}\hat{n}_{i}(\hat{n}_{i}-1) + \frac{U_{2}}{2}(\hat{\mathbf{S}}_{i}^{2}-2\hat{n}_{i}) - \mu\hat{n}_{i} - \eta\hat{S}_{iz}, \quad (1)$$

whereas the bilocal part is given by

$$\hat{H}^{(1)} = -J \sum_{\langle i,j \rangle} \sum_{\alpha} \hat{a}^{\dagger}_{i\alpha} \hat{a}_{j\alpha}.$$
(2)

Here μ and η denote the chemical potential and the external magnetic field, respectively. Furthermore, J represents the hopping matrix element between adjacent sites *i* and *j* with $\langle i, j \rangle$ indicating the summation over all nearest neighbor sites, and U_0 (U_2) stands for the on-site spin-independent (dependent) interaction strength between bosons. Additionally, $\hat{a}_{i\alpha}$ $(\hat{a}_{i\alpha}^{\dagger})$ is the annihilation (creation) operator at site *i* with hyperfine spin $\alpha \in \{-1, 0, 1\}$ which determines the total atom number operator at site *i* via $\hat{n}_i = \sum_{\alpha} \hat{a}_{i\alpha}^{\dagger} \hat{a}_{i\alpha}$ and the spin operator at site *i* according to $\hat{\mathbf{S}}_i = \sum_{\alpha,\beta} \hat{a}_{i\alpha}^{\dagger} \mathbf{F}_{\alpha\beta} \hat{a}_{i\beta}$ with the spin-1 matrices $\mathbf{F}_{\alpha\beta}$. Since the operators $\hat{\mathbf{S}}_{i}^{2}, \hat{S}_{iz}$ and \hat{n}_{i} commute with each other, their eigenvalue problems are solved by the same eigenvectors: $\hat{\mathbf{S}}_i^2 | S_i, m_i, n_i \rangle = S_i (S_i + 1) | S_i, m_i, n_i \rangle, \hat{S}_{iz} | S_i, m_i, n_i \rangle =$ $m_i |S_i, m_i, n_i\rangle$ and $\hat{n}_i |S_i, m_i, n_i\rangle = n_i |S_i, m_i, n_i\rangle$ where $S_i + n_i$ must be an even number [31, 32, 35, 36]. Thus, the eigenvalue problem of the local Hamiltonian (1) is given by

$$\hat{H}_{i}^{(0)}|S_{i}, m_{i}, n_{i}\rangle = E_{S_{i}, m_{i}, n_{i}}^{(0)}|S_{i}, m_{i}, n_{i}\rangle,$$
(3)

where the energy eigenvalues are defined as

$$E_{S_i,m_i,n_i}^{(0)} = \frac{U_0}{2} n_i (n_i - 1) + \frac{U_2}{2} \left[S_i (S_i + 1) - 2n_i \right] - \mu n_i - \eta m_i.$$
(4)

In order to artificially break the underlying U(1) symmetry of the Hamiltonian \hat{H}_{BH} , we follow [10, 11] and generalize the usual field-theoretic approach for describing classical phase transitions [37, 38] to the realm of quantum phase transitions. Thus, we couple the artificial source currents $j_{i\alpha}(\tau), j_{i\alpha}^*(\tau)$ to the operators $\hat{a}_{i\alpha}^{\dagger}$ and $\hat{a}_{i\alpha}$,

$$\hat{H}_{\rm BH}(\tau) = \hat{H}_{\rm BH} + \sum_{i} \sum_{\alpha} \left[j_{i\alpha}^*(\tau) \hat{a}_{i\alpha}(\tau) + \text{c.c.} \right], \quad (5)$$

yielding a Ginzburg–Landau theory with the spatio-temporal order parameters being defined according to

$$\Psi_{i\alpha}(\tau) = \beta \frac{\delta \mathcal{F}}{\delta j_{i\alpha}^*(\tau)}.$$
(6)

Here the free energy $\mathcal{F}[j, j^*] = -\frac{1}{\beta} \ln \mathcal{Z}[j, j^*]$ with $\beta = 1/k_{\rm B}T$ following from the partition function $\mathcal{Z}[j, j^*] = \text{Tr} \hat{T} e^{-\int_0^\beta d\tau \hat{H}_{\rm BH}(\tau)}$ with the time-ordering operator \hat{T} and the convention $\hbar = 1$. We consider equation (6) as a motivation to perform a functional Legendre transformation and define the effective action according to

$$\Gamma\left[\Psi,\Psi^*\right] = \mathcal{F}\left[j,j^*\right] - \frac{1}{\beta}\sum_i \sum_{\alpha} \left[\Psi_{i\alpha}(\tau)j_{i\alpha}^*(\tau) + \Psi_{i\alpha}^*(\tau)j_{i\alpha}(\tau)\right],$$
(7)

where $\Psi_{i\alpha}$ ($\Psi_{i\alpha}^*$) and $j_{i\alpha}^*$ ($j_{i\alpha}$) are conjugate variables satisfying the Legendre relations

$$j_{i\alpha}(\tau) = -\beta \frac{\delta\Gamma}{\delta\Psi_{i\alpha}^*(\tau)}, \qquad j_{i\alpha}^*(\tau) = -\beta \frac{\delta\Gamma}{\delta\Psi_{i\alpha}(\tau)}.$$
 (8)

In order to recover the relevant physical situation the artificial currents j^* , j should vanish. Therefore, we obtain from (8) equations of motion for determining the equilibrium value of the order parameter,

$$\frac{\delta\Gamma}{\delta\Psi_{i\alpha}^{*}(\tau)}\Big|_{\Psi=\Psi_{eq}} = 0, \qquad \frac{\delta\Gamma}{\delta\Psi_{i\alpha}(\tau)}\Big|_{\Psi=\Psi_{eq}} = 0.$$
(9)

Furthermore, we read off from equation (7) that evaluating the effective action at the equilibrium field Ψ_{eq} recovers the physical grand-canonical free energy,

$$\Gamma\left[\Psi_{\rm eq}, \Psi_{\rm eq}^*\right] = \mathcal{F}\left[0, 0\right]. \tag{10}$$

In order to calculate both the free energy \mathcal{F} and the effective action Γ , we proceed perturbatively as follows. We decompose the generalized Bose–Hubbard Hamiltonian according to $\hat{H}_{\rm BH}(\tau) = \hat{H}^{(0)} + \hat{H}^{(1)}(\tau)[j,j^*]$, where the perturbative Hamiltonian in the imaginary time Dirac interaction picture reads

$$\hat{H}_{\mathrm{I}}^{(1)}(\tau)\left[j,j^{*}\right] = -J\sum_{\langle i,j\rangle}\sum_{\alpha}\hat{a}_{i\alpha}^{\dagger}(\tau)\hat{a}_{i\alpha}(\tau) + \sum_{i}\sum_{\alpha}\sum_{\alpha}\left[j_{i\alpha}^{*}(\tau)\hat{a}_{i\alpha}(\tau) + j_{i\alpha}(\tau)\hat{a}_{i\alpha}^{\dagger}(\tau)\right].$$
(11)

With this, we determine the partition function via the Dyson series

$$\mathcal{Z} = \mathcal{Z}^{(0)} \left[1 + \sum_{n=1}^{\infty} (-1)^n \frac{1}{n!} \int_0^\beta d\tau_1 \int_0^\beta d\tau_2 \cdots \int_0^\beta d\tau_n \right] \times \left\langle \hat{T} \left[\hat{H}_{\mathrm{I}}^{(1)}(\tau_1) \hat{H}_{\mathrm{I}}^{(1)}(\tau_2) \cdots \hat{H}_{\mathrm{I}}^{(1)}(\tau_n) \right] \right\rangle^{(0)} \right]$$
(12)

with $\mathcal{Z}^{(0)} = \text{Tr } e^{-\beta \hat{H}^{(0)}}$ and the thermal average defined with respect to the unperturbed system $\langle \bullet \rangle^{(0)} = \text{Tr} [\bullet e^{-\beta \hat{H}^{(0)}}]/\mathcal{Z}^{(0)}$. The respective perturbative contributions for \mathcal{F} contain different orders of the hopping matrix element J and the currents j and j^* . As we work out a Ginzburg–Landau theory, we restrict ourselves to the fourth order in the currents. Furthermore, we focus on the leading non-trivial order in the hopping J which is of first order. Therefore, the free energy functional can be expressed in terms of imaginary time integrals over sums of products of thermal Green functions. The thermal averages in equation (12) can be expressed in terms of n-particle Green functions of the unperturbed system

$$G_{n}^{(0)}(i'_{1}\alpha'_{1},\tau'_{1};\ldots;i'_{n}\alpha'_{n},\tau'_{n}|i_{1}\alpha_{1},\tau_{1};\ldots;i_{n}\alpha_{n},\tau_{n}) = \left\langle \hat{T}\left[\hat{a}^{\dagger}_{i'_{1}\alpha'_{1}}(\tau'_{1})\hat{a}_{i_{1}\alpha_{1}}(\tau_{1})\ldots\hat{a}^{\dagger}_{i'_{n}\alpha'_{n}}(\tau'_{n})\hat{a}_{i_{n}\alpha_{n}}(\tau_{n})\right]\right\rangle^{(0)}.$$
(13)

In order to calculate the correlation functions in many-body theory, we usually use the Wick theorem which allows us to decompose the *n*-point correlation function (13) into sums of products of one-point correlation functions [37–40]. However, this theorem is not valid for the considered system here because the unperturbed Bose–Hubbard Hamiltonian (1) contains terms which are of fourth order in the creation and annihilation operators. Therefore, we use the cumulant decomposition for the Green functions, which is based on the locality of $\hat{H}^{(0)}$ [41, 42]. With this, the unperturbed one- and two-point Green functions are given by

$$G_1^{(0)}(i_1\alpha_1, \tau_1 | i_2\alpha_2, \tau_2) = \delta_{i_1, i_2 i_1} C_1^{(0)}(\tau_1, \alpha_1 | \tau_2, \alpha_2), \quad (14)$$

and

$$G_{2}^{(0)}(i_{1}\alpha_{1}, \tau_{1}; i_{2}\alpha_{2}, \tau_{2}|i_{3}\alpha_{3}, \tau_{3}; i_{4}\alpha_{4}, \tau_{4})$$

$$= \delta_{i_{1},i_{3}}\delta_{i_{2},i_{4}}\delta_{i_{3},i_{4}i_{1}}C_{1}^{(0)}(\tau_{1}, \alpha_{1}; \tau_{2}, \alpha_{2}|\tau_{3}, \alpha_{3}; \tau_{4}, \alpha_{4})$$

$$+ \delta_{i_{1},i_{3}}\delta_{i_{2},i_{4}i_{1}}C_{1}^{(0)}(\tau_{1}, \alpha_{1}|\tau_{3}, \alpha_{3})_{i_{1}}C_{1}^{(0)}(\tau_{2}, \alpha_{2}|\tau_{4}, \alpha_{4})$$

$$+ \delta_{i_{1},i_{4}}\delta_{i_{2},i_{3}i_{1}}C_{1}^{(0)}(\tau_{1}, \alpha_{1}|\tau_{4}, \alpha_{4})_{i_{1}}C_{1}^{(0)}(\tau_{2}, \alpha_{2}|\tau_{3}, \alpha_{3}).$$
(15)

In order to calculate the respective cumulants from combining equations (13)–(15), we use for each lattice site the property [32, 43, 44]

$$\hat{a}_{\alpha}^{\dagger}|S,m,n\rangle = M_{\alpha,S,m,n}|S+1,m+\alpha,n+1\rangle + N_{\alpha,S,m,n}|S-1,m+\alpha,n+1\rangle,$$
(16)

$$\hat{a}_{\alpha}|S,m,n\rangle = O_{\alpha,S,m,n}|S+1,m-\alpha,n-1\rangle + P_{\alpha,S,m,n}|S-1,m-\alpha,n-1\rangle,$$
(17)

where $M_{\alpha,S,m,n}$, $N_{\alpha,S,m,n}$, $O_{\alpha,S,m,n}$ and $P_{\alpha,S,m,n}$ represent recursively defined matrix elements of creation and annihilation operators. Having calculated the free energy \mathcal{F} in this way, we perform then the Legendre transformation (7) to determine the effective action. In the special case of a stationary equilibrium, which is site-independent due to homogeneity, the order parameter is given in terms of Matsubara frequencies $\omega_m = 2\pi m/\beta$: $\Psi_{i\alpha}^{eq}(\omega_m) = \Psi_{\alpha}\sqrt{\beta} \,\delta_{m,0}, \Psi_{i\alpha}^{eeq}(\omega_m) = \Psi_{\alpha}^*\sqrt{\beta} \,\delta_{m,0}$. Thus, the on-site effective potential becomes

$$\Gamma\left(\Psi_{\alpha}, \Psi_{\alpha}^{*}\right) = \mathcal{F}_{0} + \sum_{\alpha} B_{\alpha} |\Psi_{\alpha}|^{2} + \sum_{\alpha_{1},\alpha_{2},\alpha_{3},\alpha_{4}} A_{\alpha_{1}\alpha_{2}\alpha_{3}\alpha_{4}} \Psi_{\alpha_{1}}^{*} \Psi_{\alpha_{2}}^{*} \Psi_{\alpha_{3}} \Psi_{\alpha_{4}}, \qquad (18)$$

with the Landau coefficients

$$B_{\alpha} = \frac{1}{a_2^{(0)}(\alpha, 0)} - zJ,$$
(19)

$$A_{\alpha_1\alpha_2\alpha_3\alpha_4} = -\frac{\beta a_4^{(0)}(\alpha_1, 0; \alpha_2, 0 | \alpha_3, 0; \alpha_4, 0)}{4 a_2^{(0)}(\alpha_1, 0) a_2^{(0)}(\alpha_2, 0) a_2^{(0)}(\alpha_3, 0) a_2^{(0)}(\alpha_4, 0)},$$
(20)

where z = 2D denotes the coordination number in a *D*-dimensional cubic lattice [45]. Furthermore, $a_2^{(0)}$ and $a_4^{(0)}$ follow from the cumulants but they are not displayed here due to their complicated and lengthy expressions.

Extremizing the effective potential (18) according to (9) we find at first the location of the quantum phase transition

$$zJ_{\rm c} = \frac{\min}{\alpha} \frac{1}{a_2^{(0)}(\alpha, 0)},$$
 (21)

which turns out to coincide with the mean-field result in [32–34], see figure 1. Moreover, inserting (18) into (9) yields also the different superfluid phases for ferromagnetic and anti-ferromagnetic interactions with and without magnetization at zero temperature. If there is more than one solution, we must take the one which minimizes the effective potential for some system parameter. At first, we observe that the condensate in the superfluid phase above the first Mott lobe shows, indeed, a sharp increase [11]. Thus, the condensate density cannot be valid deep in the superfluid phase. Therefore, it is necessary to determine the range of validity of the Ginzburg–Landau theory. To this end, we use the fact that we cannot have more particles in the condensate than we have in total. This leads to the condition that the sum over the condensate densities $\sum_{\alpha} |\Psi_{\alpha}|^2$ is equal to the average number of particles per lattice site $\langle n \rangle = -\frac{\partial \Gamma}{\partial \mu}|_{\Psi=\Psi_{eq}}$, i.e.

$$\sum_{\alpha} |\Psi_{\alpha}|^2 = \langle n \rangle, \qquad (22)$$

which is graphically shown in figure 1. However, we read off from figure 1 that condition (22) breaks down at the end of the lower Mott lobes. There we have to use an additional criterion to obtain a finite range of validity. To this end we complement condition (22) by the additional ad hoc restriction that above



Figure 1. Superfluid phases calculated analytically and numerically for different values of spin-dependent interaction strength U_2/U_0 and at external field $\eta/U_0 = 0.2$: $\Psi_1 \neq 0$, $\Psi_0 = \Psi_{-1} = 0$ (blue); $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 \neq 0$ (red); $\Psi_{-1} \neq 0$, $\Psi_0 = \Psi_1 = 0$ (cyan); and $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 = 0$ (green). The validity ranges (22) and (23) correspond to the black and dashed orange lines, respectively. Moreover, Mott lobes are characterized by gray color. (a) $U_2/U_0 = 0.02$. (b) $U_2/U_0 = 0.04$. (c) $U_2/U_0 = 0.05$. (d) $U_2/U_0 = 0.07$. (e) $U_2/U_0 = 0.15$.

Mott lobe *n* the condensate density cannot be larger than n+1, yielding the boundary

$$\sum_{\alpha} |\Psi_{\alpha}|^2 = n+1, \tag{23}$$

which is depicted in figure 1 as a dashed orange line.

Now we show at zero temperature that our Ginzburg– Landau theory distinguishes between various superfluid phases for a ferromagnetic and anti-ferromagnetic interaction with and without external magnetic field within the validity range of our theory. Without external magnetization the superfluid phase is a polar (ferromagnetic) state for anti-ferromagnetic (ferromagnetic) interactions, which is characterized by $\Psi_1 \neq 0$, $\Psi_{-1} = \Psi_0 = 0$ ($\Psi_0 \neq 0$, $\Psi_{-1} =$ $\Psi_1 = 0$), in accordance with previous mean-field results [31, 32]. In the presence of the magnetic field the phase diagram does not change for the ferromagnetic interaction, as the minimization of the energy implies the maximum of spin value as it is in the case without η , except the degeneracy with respect to *m* is lifted, so the ground state becomes $|n, n, n\rangle$. For an anti-ferromagnetic interaction, however, the situation is more complicated with an external magnetic field due to two competing effects. Whereas the anti-ferromagnetic interaction energetically favors anti-parallel spins, the external magnetic field has the tendency to align the spins. In the following we show that this competition leads to the appearance of different superfluid phases with different magnetic properties.

In figure 1, we study the predictions of the Ginzburg– Landau theory on how the external magnetic field affects the superfluid phases in the case of an anti-ferromagnetic interaction, i.e. $U_2 > 0$. In this context it is important to determine the degeneracy when two states have the same energy with equal particle number but different total spin. Using the ground state energy (4) and the degeneracy condition $E_{S,S,n}^{(0)} = E_{S+2,S+2,n}^{(0)}$, we obtain the critical spin-dependent interaction strength $U_2^{\text{crit}} = \eta/(S + \frac{3}{2})$ at the external magnetic field η [34, 43, 44]. With this, we get the resulting phase diagrams below and above the critical spin-dependent interaction strength.

When U_2/U_0 is 0.02, the superfluid phase becomes $\Psi_1 \neq 0$, $\Psi_{-1} = \Psi_0 = 0$ where the ground state $|n, n, n\rangle$ is the state with maximum spin for all six lobes as shown in figure 1(a). Above the first critical value $U_{2\text{even}}^{(1)}/U_0 = 0.036$ both the spin *S* and the magnetic *m* quantum numbers change from $|6, 6, n\rangle$ to $|4, 4, n\rangle$ for even lobes as shown in figure 1(b). We remark that if the spin-dependent interaction increases, the effect of the external magnetic field decreases, so the Mott lobes increase. The phases $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 = 0$ and $\Psi_{-1} \neq 0$, $\Psi_1 = \Psi_0 = 0$ appear in the SF phase for the fifth and sixth lobes. We note that the phase $\Psi_{-1} \neq 0$, $\Psi_1 \neq 0$, $\Psi_0 = 0$ appears twice in the sixth lobe. The right phase in the sixth lobe from $|7, 7, 7\rangle$ to $|5, 5, 7\rangle$, which happens at the critical value $U_{2odd}^{(1)}/U_0 = 0.0308$.

Beyond the critical value $U_{2odd}^{(2)}/U_0 = 0.044$ the values of *S* and *m* for the odd lobes change from $|5, 5, n\rangle$ to $|3, 3, n\rangle$ as shown in figure 1(c). Similarly, the phases $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 = 0$ and $\Psi_{-1} \neq 0$, $\Psi_1 = \Psi_0 = 0$ appear in the SF phase for the fourth and fifth lobe and the phase $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 = 0$ increases in the sixth lobe. After the critical value $U_{2even}^{(2)}/U_0 = 0.05714$, the ground states for the even lobes change from $|4, 4, n\rangle$ to $|2, 2, n\rangle$ as shown in figure 1(d). By the same way the phases $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 = 0$ and $\Psi_{-1} \neq 0$, $\Psi_1 = \Psi_0 = 0$ are seen in the SF phase for the third and fourth lobe.

When U_2 increases beyond the critical value $U_{2odd}^{(3)}/U_0 = 0.08$, the ground states for the odd lobes change from $|3, 3, n\rangle$ to $|1, 1, n\rangle$ as shown in figure 1(e). The phases $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 = 0$ and $\Psi_{-1} \neq 0$, $\Psi_1 = \Psi_0 = 0$ appear in the SF phase for the even and odd lobes. After the critical value $U_{2even}^{(3)}/U_0 = 0.133$, S and m change from $|2, 2, n\rangle$ to $|0, 0, n\rangle$ for the even lobes as shown in figure 1(f). Furthermore, the effect of magnetic field becomes very weak because the value of η is close to U_2 . Additionally, spin pairs are produced to get the minimal energy and, thus, the ground state becomes

 $|0, 0, n\rangle$ for an even *n*, and $|1, 1, n\rangle$ for an odd *n*. We found the new phase $\Psi_{-1} \neq 0$, $\Psi_1 \neq 0$, $\Psi_0 \neq 0$ above the even lobes.

Furthermore, inspecting the energies of the respective phases in the vicinity of their boundaries allows us to determine the order of the quantum phase transition. With this we find that the quantum phase transition from the Mott insulator to the superfluid phase is of second order for spin-1 bosons in a cubic optical lattice under the effect of the magnetic field at zero temperature. Thus, our finding disagrees with Kimura et al [46] where a first-order SF-MI phase transition was found at a part of the phase boundary by using the Gutzwiller variational approach. Finally, we observe that the transitions between the different superfluid phases can be of both first and second order above the same Mott lobe. For instance, the transition from $\Psi_1 \neq 0$, $\Psi_0 = \Psi_{-1} = 0$ to $\Psi_{-1} \neq 0, \Psi_1 = \Psi_0 = 0$ or vice versa is of first order, whereas the transition from $\Psi_1 \neq 0$, $\Psi_0 = \Psi_{-1} = 0$ to $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$ 0, $\Psi_0 = 0$ or $\Psi_1 \neq 0$, $\Psi_{-1} \neq 0$, $\Psi_0 \neq 0$ phases or vice versa is of second order.

In conclusion, we have worked out a Ginzburg–Landau theory for spin-1 bosons in a cubic optical lattice within its range of validity and investigated analytically and numerically at zero temperature the different superfluid phases for an anti-ferromagnetic interaction in the presence of an external magnetic field. Depending on the particle number, the spin-dependent interaction and the value of the magnetic field, we find superfluid phases with a macroscopic occupation of the two spin states ± 1 or even of all three spin states $0, \pm 1$. This is different from the mean-field approximation which only predicted two superfluid phases with spins aligned or opposite to the field direction [33, 34]. It would be interesting to study how these results would change in a frustrated triangular optical lattice [30] or in a superlattice [47].

Acknowledgments

We acknowledge discussions with Antun Balaz and Mathias Ohliger as well as financial support from both the Egyptian Government and the German Research Foundation (DFG) via the Collaborative Research Center SFB/TR49 Condensed Matter Systems with Variable Many-Body Interactions.

References

- [1] Yukalov V I 2009 Laser Phys. 19 1
- [2] Lewenstein M, Sanpera A and Ahufinger V 2012 Ultracold Atoms in Optical Lattices, Simulating Quantum Many-Body Systems (Oxford: Oxford University Press)
- [3] Greiner M, Mandel O, Esslinger T, Hänsch T W and Bloch I 2002 Nature 415 39
- [4] Greiner M, Mandel O, Hänsch T W and Bloch I 2002 Nature 419 51
- [5] Fisher M P A, Weichman P B, Grinstein G and Fisher D S 1989 Phys. Rev. B 40 546
- [6] Jaksch D, Bruder C, Cirac J I, Gardiner C W and Zoller P 1998 Phys. Rev. Lett. 81 3108
- [7] Sachdev S 2001 *Quantum Phase Transitions* 2nd edn (Cambridge: Cambridge University Press)
- [8] Freericks J K and Monien H 1996 Phys. Rev. B 53 2691
- [9] van Oosten D, Straten P and Stoof H T C 2001 Phys. Rev. A 63 053601

- [10] dos Santos F E A and Pelster A 2009 Phys. Rev. A 79 013614
- [11] Bradlyn B, dos Santos F E A and Pelster A 2009 Phys. Rev. A
- 79 013615
 [12] Teichmann N, Hinrichs D, Holthaus M and Eckardt A 2009 *Phys. Rev. B* 79 100503
- [13] Teichmann N, Hinrichs D, Holthaus M and Eckardt A 2009 Phys. Rev. B 79 224515
- [14] Hinrichs D, Pelster A and Holthaus M 2013 Appl. Phys. B at press
- [15] Capogrosso-Sansone B, Prokof'ev N V and Svistunov B V 2007 Phys. Rev. B 75 134302
- [16] Capogrosso-Sansone B, Söyler S G, Prokof'ev N and Svistunov B 2008 Phys. Rev. A 77 015602
- [17] Wang T, Zhang X-F, Eggert S and Pelster A 2013 Phys. Rev. A 87 063615
- [18] Modugno G, Ferlaino F, Heidemann R, Roati G and Inguscio M 2003 Phys. Rev. A 68 011601(R)
- [19] Albus A, Illuminati F and Eisert J 2003 Phys. Rev. A 68 023606
- [20] Büchler H P and Blatter G 2003 Phys. Rev. Lett. 91 130404
- [21] Lewenstein M, Santos L, Baranov M and Fehrmann H 2004 Phys. Rev. Lett. 92 050401
- [22] Bloch I 2008 Nature **453** 1016
- [23] Krutitsky K V, Pelster A and Graham R 2006 New J. Phys. 8 187
- [24] Roati G, D'Errico C, Fallani L, Fattori M, Fort C, Zaccanti M, Modugno G, Modugno M and Inguscio M 2008 Nature 453 895
- [25] Bissbort U, Thomale R and Hofstetter W 2010 Phys. Rev. A 81 063643
- [26] Stenger J, Inouye S, Stamper-Kurn D M, Miesner H-J, Chikkatur A P and Ketterle W 1999 Nature 396 345
- [27] Barrett M, Sauer J and Chapman M S 2001 Phys. Rev. Lett. 87 010404

- [28] Widera A, Gerbier F, Fölling S, Gericke T, Mandel O and Bloch I 2005 Phys. Rev. Lett. 95 190405
- [29] Widera A, Gerbier F, Fölling S, Gericke T, Mandel O and Bloch I 2006 New J. Phys. 8 152
- [30] Becker C, Soltan-Panahi P, Kronjäger J, Dörscher S, Bongs K and Sengstock K 2010 New J. Phys. 12 065025
- [31] Demler E and Zhou F 2002 Phys. Rev. Lett. 88 163001
- [32] Tsuchiya S, Kurihara S and Kimura T 2004 Phys. Rev. A 70 043628
- [33] Svidzinsky A A and Chui S T 2003 Phys. Rev. A 68 043612
- [34] Uesugi N and Wadati M 2003 J. Phys. Soc. Japan 72 1041
- [35] Wu Y 1996 Phys. Rev. A 54 4534
- [36] Kawaguchi Y and Ueda M 2012 *Phys. Rep.* **520** 253
- [37] Kleinert H and Schulte-Frohlinde V 2001 Critical Properties of Φ⁴-Theories (Singapore: World Scientific)
- [38] Zinn-Justin J 2002 Quantum Field Theory and Critical Phenomena 4th edn (Oxford: Claredon)
- [39] Gorter C J and Casimir H G B 1934 Phys. Z. 35 963
- [40] Kadanoff L P and Baym G 1962 Quantum Statistical Mechanics: Green's Function Methods in Equilibrium and Non-Equilibrium Problems (New York: Benjamin)
- [41] Metzner W 1991 Phys. Rev. B 43 8549
- [42] Ohliger M and Pelster A 2013 World J. Condens. Matter Phys. 3 125
- [43] Kimura T, Tsuchiya S, Yamashita M and Kurihara S 2006 J. Phys. Soc. Japan 75 074601
- [44] Ohliger M 2008 Thermodynamic properties of spinor bosons in optical lattices *Diploma Thesis* Freie Universität Berlin
- [45] Mobarak M, Spinor Bose gases in cubic optical lattice *PhD Thesis* Freie Universität Berlin, in preparation
- [46] Kimura T, Tsuchiya S and Kurihara S 2005 Phys. Rev. Lett. 94 110403
- [47] Wagner A, Nunnenkamp A and Bruder C 2012 Phys. Rev. A 86 023624